

On the Number of Graphical Forest Partitions

Deborah A. Frank, Carla D. Savage and James A. Sellers

Department of Mathematics
Miami University, Hamilton
1601 Peck Boulevard
Hamilton, OH 40511

Department of Computer Science, Box 8206
North Carolina State University
Raleigh, NC 27695

Department of Science and Mathematics
Cedarville University
Cedarville, OH 45314

February 6, 2001

Abstract

A **graphical partition** of the even integer n is a partition of n where each part of the partition is the degree of a vertex in a simple graph and the degree sum of the graph is n . In this note, we consider the problem of enumerating a subset of these partitions, known as **graphical forest partitions**, graphical partitions whose parts are the degrees of the vertices of forests (disjoint unions of trees). We shall prove that

$$gf(2k) = p(0) + p(1) + p(2) + \dots + p(k-1)$$

where $gf(2k)$ is the number of graphical forest partitions of $2k$ and $p(j)$ is the ordinary partition function which counts the number of integer partitions of j .

1 Introduction

A **partition** of a positive integer n is a sequence of positive integers, in no particular order, whose sum is n . For example, $5 + 3 + 2 + 2 + 1 + 1 + 1$ is a partition of 15. Each number in a partition is called a **part** of that partition. The partition function $p(n)$ counts the number of partitions of the integer n .

In this note we will consider only those partitions of n that are graphical sequences (and denote the number of these partitions by $g(n)$). A **graphical sequence** is a sequence whose terms represent the degrees of the vertices in a **simple graph**, a graph that can be drawn without any multiple edges or loops.

Finding a closed formula for $g(n)$ has proven difficult. Indeed, even the asymptotics of $g(n)$ are still unknown. However, several results regarding $g(n)$ are known. For instance, a lower bound for this function has been found. This lower bound is $p(n) - p(n-1)$, which is also the number of partitions of n with all successive ranks negative [4]. Moreover, it is also known [7] that an upper bound for $g(n)$ is $(.25 + o(1))p(n)$.

Finally, Pittel [6] has shown that

$$\frac{g(n)}{p(n)} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

The interested reader may also wish to see [2] and [5] for additional discussion regarding $g(n)$.

Because of the difficulty in finding a closed formula for $g(n)$, we chose to restrict $g(n)$ even further by considering only those graphical partitions of n which correspond to forests, with the hope that a closed form might become apparent. (Here we use the term **forest** to mean a union of trees.) We denote the number of graphical forest partitions of n by $gf(n)$.

The goal of this note is to prove that, for all $k \geq 1$,

$$gf(2k) = p(0) + p(1) + p(2) + \dots + p(k-1).$$

2 The Results

First off, we let $gf(n, t)$ be the number of graphical forest partitions of n into exactly t parts. Our first goal is to prove the following result.

Theorem 2.1. *For $s > 1$, $gf(2k, k + s) = gf(2k - 2, k + s - 2)$.*

Proof. If a forest realizes a sequence counted by $gf(2k, k + s)$, at least two of its vertices in different components must have degree 1. Deleting these two and joining their neighbors by an edge gives a forest realization of a sequence counted by $gf(2k - 2, k + s - 2)$. Conversely, adding two new vertices joined by an edge to a forest realization of a sequence counted in $gf(2k - 2, k + s - 2)$ creates a new sequence counted by $gf(2k, k + s)$. \square

We turn now to our main theorem.

Theorem 2.2. *For all $k \geq 2$, $gf(2k) = p(0) + p(1) + \dots + p(k - 1)$.*

Proof. It is known [8, Problem 2.1.12] that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{k+1} > 0$ is the degree sequence of a tree if and only if $\lambda_1 + \lambda_2 + \dots + \lambda_{k+1} = 2k$. Letting $p(n, t)$ denote the number of partitions of n with exactly t parts, then the number of graphical tree partitions of $2k$ is $p(2k, k + 1)$ which equals $p(k - 1)$.

Thus, $gf(2k, k + 1) = p(k - 1)$ and

$$gf(2k) = \sum_{s=1}^k gf(2k, k + s).$$

Finally, from Theorem 2.1 above, we know $gf(2k, k + s) = gf(2k - 2, k + s - 2)$ and the result follows. \square

3 On Computing $gf(2k)$

Thanks to the results above, we see that finding the number of graphical forest partitions of $2k$ simply involves finding the values of the ordinary partition function $p(n)$, $0 \leq n \leq k - 1$.

A quick word on asymptotics is worth noting here. It is known [3, Section 3] that

$$p(0) + \dots + p(n - 1) \sim p(n) \frac{\sqrt{6n}}{\pi},$$

which means

$$gf(2k) \sim p(k) \frac{\sqrt{6k}}{\pi}.$$

Moreover, we know [1, p. 70] that

$$p(k) \sim \frac{1}{4k\sqrt{3}} \exp \left[\pi \left(\frac{2k}{3} \right)^{1/2} \right],$$

which implies

$$gf(2k) \sim \frac{\sqrt{2}}{4\pi\sqrt{k}} \exp \left[\pi \left(\frac{2k}{3} \right)^{1/2} \right].$$

Next, we mention two ways to determine exact values of $gf(2k)$. First, we can utilize the generating function for $p(n)$ and Euler's Pentagonal Number Theorem to develop the following recurrence for $p(n)$:

$$p(n) = p(n-1) + p(n-2) - p(n-5) - p(n-7) + p(n-12) + p(n-15) - \dots$$

where the values being subtracted in the arguments on the right-hand side are the pentagonal numbers $\frac{3}{2}m^2 - \frac{1}{2}m$ for integers m . The interested reader should see [1, p. 11].

Alternatively, we can compute the values of $gf(2k)$ by developing a generating function for $gf(2k)$ and expanding it using a computer algebra system. Since the generating function for $p(n)$ is given by

$$\sum_{k=0}^{\infty} p(k)q^k = \prod_{n \geq 1} \frac{1}{1-q^n},$$

we see that the coefficient of q^k in

$$\frac{q}{1-q} \prod_{n \geq 1} \frac{1}{1-q^n}$$

is $p(0) + p(1) + \dots + p(k-1)$. Thus,

$$\sum_{k=0}^{\infty} gf(2k)q^k = \frac{q}{1-q} \prod_{n \geq 1} \frac{1}{1-q^n}.$$

References

- [1] G. E. Andrews, "Theory of Partitions," Addison-Wesley, 1976.
- [2] T. M. Barnes and C. D. Savage, Efficient generation of graphical partitions, *Discrete Applied Mathematics*, **78** no. 1-3 (1997), 17-26.
- [3] S. Corteel, B. Pittel, C. D. Savage, and H. Wilf, On the multiplicity of parts in a random partition, *Random Structures and Algorithms*, **14**, no. 2 (1999) 185-197.
- [4] P. Erdős and L. B. Richmond, On graphical partitions, *Combinatorica*, **13**, no. 1 (1993), 57-63.
- [5] J. M. Nolan, V. Sivaraman, C. D. Savage, and P. K. Tiwari, Graphical basis partitions, *Graphs and Combinatorics*, **14**, no. 3 (1998), 241-261.
- [6] B. Pittel, Confirming two conjectures about the integer partitions. *J. Combinatorial Theory Series A*, **88**, no. 1 (1999), 123-135.

- [7] C. C. Rousseau and F. Ali, A note on graphical partitions, *Journal of Combinatorial Theory Series B*, **64** (1995), 314–318.
- [8] D. West, “Introduction to Graph Theory,” First Edition, Prentice Hall, 1995.

2000 *Mathematics Subject Classification.* 05A17